

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
31 December 2003 (31.12.2003)

PCT

(10) International Publication Number
WO 2004/000829 A1

(51) International Patent Classification⁷: **C07D 401/14**,
401/12, 403/04, A61K 31/444, 31/497, A61P 25/00

S-753 28 Uppsala (SE). ANDERSSON, Emma; Smed-
vägen 14, S-176 71 Järfälla (SE). NYGREN, Alf;
Ingvarsgratan 47, S-753 34 Uppsala (SE).

(21) International Application Number:
PCT/SE2003/001043

(74) Agent: CONRADI, Magnus; Biovitrum AB, Lindhagens-
gatan 133, S-112 76 Stockholm (SE).

(22) International Filing Date: 18 June 2003 (18.06.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

0201881-0	19 June 2002 (19.06.2002)	SE
60/390,656	21 June 2002 (21.06.2002)	US
0202041-0	28 June 2002 (28.06.2002)	SE
0202516-1	26 August 2002 (26.08.2002)	SE
60/406,119	26 August 2002 (26.08.2002)	US
60/416,701	7 October 2002 (07.10.2002)	US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: BIOVITRUM AB [SE/SE]; Lindhagensgatan 133, S-112 76 Stockholm (SE).


(72) Inventors: CERNERUD, Magnus; Nybrogatan 68, S-114 41 Stockholm (SE). LUNDSTRÖM, Helena; Mjölnerbacken 15, 1 trp, S-174 48 Sundbyberg (SE). LÖFSTRÖM, Claes; Stadshagsplan 10, S-112 50 Stockholm (SE). PELCMAN, Margit; Industrigatan 1, S-112 46 Stockholm (SE). FLECK, Tom (deceased). PAPTCHIKHINE, Alexander; Petterslundsgatan 15,

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL PROCESS



(57) Abstract: The present invention relates to a process for the preparation of compounds which are therapeutically active in the central nervous system. In one aspect, the invention relates to a process for the preparation of compounds of the general formula (I). The invention also relates to (2R)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine, hydrochloride and a process for preparing the same. Another object of the present invention is a compound of the formula (I). Another object of the present invention is a method for the treatment or prophylaxis of a serotonin-related disorder, comprising administering to a subject in need thereof an effective amount of the compound above; as well as the use of said compound to manufacture a medicament for the treatment of a serotonin-related disorder.

NOVEL PROCESS

Related applications

This application claims priority to Swedish application number 0201881-0, filed on
5 June 19, 2002, Swedish application number 0202041-0, filed on June 28, 2002, Swedish
application number 0202516-1, filed on August 26, 2002, U.S. provisional application
60/390,656, filed June 21, 2002, U.S. provisional application 60/406,119, filed on August 26,
2002, and U.S. provisional application 60/416,701, filed on October 7, 2002, the contents of
which are incorporated herein by reference.

10

Field of the invention

The present invention relates to a process for the preparation of compounds which
are therapeutically active in the central nervous system.

15

Background of the invention

Many diseases of the central nervous system are influenced by the adrenergic, the
dopaminergic, and the serotonergic neurotransmitter systems. For example, serotonin has
been implicated in a number of diseases and conditions which originate in the central nervous
system. A number of pharmacological and genetic experiments involving receptors for
20 serotonin strongly implicate the 5-HT_{2C} receptor subtype in the regulation of food intake
(Obes. Res. 1995, 3, Suppl. 4, 449S-462S). The 5-HT_{2C} receptor subtype is transcribed and
expressed in hypothalamic structures associated with appetite regulation. It has been
demonstrated that the non-specific 5-HT_{2C} receptor agonist *m*-chlorophenylpiperazine
(mCPP), which has some preference for the 5-HT_{2C} receptor, causes weight loss in mice that
25 express the normal 5-HT_{2C} receptor while the compound lacks activity in mice expressing the
mutated inactive form of the 5-HT_{2C} receptor (Nature 1995, 374, 542-546). In a recent
clinical study, a slight but sustained reduction in body weight was obtained after 2 weeks of
treatment with mCPP in obese subjects (Psychopharmacology 1997, 133, 309-312). Weight
reduction has also been reported from clinical studies with other "serotonergic" agents (see
30 e.g. IDrugs 1998, 1, 456-470). For example, the 5-HT reuptake inhibitor fluoxetine and the 5-
HT releasing agent/reuptake inhibitor dexfenfluramine have exhibited weight reduction in
controlled studies. However, currently available drugs that increase serotonergic transmission
appear to have only a moderate and, in some cases, transient effects on the body weight.

The 5-HT_{2c} receptor subtype has also been suggested to be involved in CNS disorders such as depression and anxiety (Exp. Opin. Invest. Drugs 1998, 7, 1587-1599; IDrugs, 1999, 2, 109-120).

The 5-HT_{2c} receptor subtype has further been suggested to be involved in urinary disorders such as urinary incontinence (IDrugs, 1999, 2, 109-120).

Compounds which have a selective effect on the 5-HT_{2c} receptor may therefore have a therapeutic potential in the treatment or prophylaxis of disorders like those mentioned above. Of course, selectivity also reduces the potential for adverse effects mediated by other serotonin receptors.

Examples of such compounds are (2*R*)-1-(3-{2-[(2-ethoxy-3-pyridinyl)oxy]ethoxy}-2-pyrazinyl)-2-methylpiperazine, (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine and pharmaceutically acceptable acid salts thereof. WO 00/76984 (hereinafter called D1) relates to a process for the preparation of such compounds on a small scale such as a gram scale. A problem to be solved by the present invention was to prepare such compounds on a large scale such as on a kilogram scale. The following factors are more important for preparation on a large scale, in comparison to preparation on a small scale:

- to obtain a high yield of the desired products for economy reasons,
- that the processes for preparation are safe with regard to explosion,
- that the reagents and solvents used are relatively non-toxic,
- that the products obtained are relatively stable, and
- that the reaction times are relatively short.

These problems have been solved by the present invention. It has been shown that the yields of the desired products according to the present invention are higher than the yields according to D1. In the experimental part, the yields according to the present invention and D1 are compared. Regarding the choice of solvents for the process steps, dioxane, as used according to D1, has been replaced by solvents such as MtBE and THF (see step (ii) below) which are less prone to form peroxides and which are less carcinogenic than dioxane. Furthermore, it has been shown that (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, L-malate salt prepared according to the present invention (see Example 3A) has superior properties compared to (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, hydrochloride prepared according to D1 in that the former has a higher crystallinity, is less hygroscopic and has a higher chemical stability than the latter. Regarding chemical stability, D1 discloses the preparation of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine, which has

also been prepared in Example 8 below. This compound is not stable for storage as a free base. As (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine is a key intermediate, chemical stability during long term storage is important with regard to process economy. It has now been found that the corresponding hydrochloride salt thereof is considerably more stable,
5 which has been prepared in Example 9 below.


The method to prepare Example 3C is a good way of increasing the purity of Example 2C. It has been shown that Example 2C with a purity of 60-70% gives Example 3C with a purity of 99% in one crystallization step. By contrast, the same purity increasing effect has not been achieved by making the acetate of Example 2C.

10 Regarding reaction time, the reaction according to Example 2A below was complete at room temperature in 15 minutes. The same compound has been prepared in Example 173 in D1. The procedure of Example 172, step 3 has been followed, wherein the reaction was stirred at 85°C for 15 h. Furthermore, the reaction according to Example 2B below was complete in 35 minutes at 55°C. The same compound has been prepared in Example 200 in
15 D1. The procedure of Example 192, step 3 has been followed, wherein the reaction was stirred at 90°C for 2 h.

Summary of the invention

One object of the present invention is a process for the preparation of compounds
20 which bind to the 5-HT_{2C} receptor (agonists and antagonists) and which therefore may be used for the treatment or prophylaxis of serotonin-related disorders.


In one aspect, the invention relates to a process for the preparation of compounds of the general formula (I):




comprising the steps of:

- (i) hydroxyalkylating (i.e., O-alkylating) a 3-pyridinol derivative of the general formula (II) or the corresponding hydrochloride:

5




to give another 3-pyridinol derivative of the general formula (III):




- 10 (ii) condensing the 3-pyridinol derivative of the general formula (III) with (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpyrazine of the formula (IV) in the presence of an alkali metal tert-butoxide (e.g. potassium tert-butoxide) or an alkali earth metal tert-butoxide, as a base in a solvent system (e.g., a solvent system containing tetrahydrofuran),

15



to give a compound of the general formula (V):




- 5 (iii) which is then converted to the compound of the general formula (I) by treatment with a pharmaceutically acceptable acid of the formula HA,

wherein


- 10 R₁-R₄ are each independently selected from the group consisting of hydrogen, halogen, C₁-C₆-alkyl, C₁-C₆-alkoxy, and di-C₁-C₆-alkylamino-C₂-C₆-alkoxy; and wherein

(2R)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) is prepared by:

- (iv) acidification of racemic 2-methylpiperazine of the formula (VI):




with L-tartaric acid and fractional crystallization to give (*R*)-2-methylpiperazine, L-tartrate of the formula (VII):




5


- (v) basification of (*2R*)-2-methylpiperazine, L-tartrate of the formula (VII) to give (*R*)-2-methylpiperazine of the formula (VIII):



- 10 (vi) tritylation of (*R*)-2-methylpiperazine of the formula (VIII) to give (*R*)-3-methyl-1-tritylpiperazine of the formula (IX):



- (vii) condensation of (*R*)-3-methyl-1-tritylpiperazine of the formula (IX) with 2,3-dichloropyrazine to give (*2R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X):



5

- (viii) detritylation of (*2R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X) to give (*2R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV),
- (ix) and conversion of (*2R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) to a suitable acid addition salt (e.g. hydrochloride salt).

10

It is preferred that R_1 - R_3 are hydrogen and R_4 is selected from the group consisting of hydrogen, ethoxy, and 2-dimethylaminoethoxy.

It is also preferred that:

- 15 - step (iv) is performed by using water and ethanol as solvents;
- step (v) is performed by using a hydroxide (e.g., sodium hydroxide) as a base;
- step (vi) is performed with trityl chloride in the presence of triethylamine;
- step (vii) is performed in the presence of an alkali metal carbonate (e.g. potassium carbonate) or an alkali earth metal carbonate, with dimethyl formamide as a solvent; and
- 20 - step (viii) is performed in 10% sulfuric acid in isopropanol.

When R_1 - R_4 are hydrogen, it is preferred that:

- step (i) is performed by reacting the compound of the formula (II) with ethylene carbonate with an alkali metal carbonate (e.g. potassium carbonate) or an alkali earth metal carbonate, as a base and dimethyl formamide as a solvent,

25

- step (ii) is performed by reacting the compound of the formula (III) with the compound of the formula (IV) in the presence of an alkali metal tert-butoxide (e.g. potassium tert-butoxide) or an alkali earth metal tert-butoxide, in tetrahydrofuran,
- the pharmaceutically acceptable acid of the formula HA in step (iii) is L-malic acid.

5

Before converting the compound of the formula (I), wherein R_1 - R_4 are hydrogen, into the L-malate salt, the product of the formula (V) is preferably converted to the fumarate, giving a more pure product. After that, the fumarate may be converted to the L-malate, e.g. via neutralization of the fumarate with a base and then converted to the L-malate by means of L-malic acid.

10

When R_1 - R_3 are hydrogen and R_4 is ethoxy, it is preferred that:

- step (i) is performed by reacting the hydrochloride of the compound of the formula (II) with 2-chloroethanol with in an aqueous solution of a hydroxide, more preferably sodium hydroxide,
- step (ii) is performed by reacting the compound of the formula (III) with the compound of the formula (IV) in the presence of an alkali metal tert-butoxide (e.g. potassium tert-butoxide) or an alkali earth metal tert-butoxide, in a solvent system consisting of methyl tert-butylether and tetrahydrofuran,
- the pharmaceutically acceptable acid of the formula HA in step (iii) is succinic acid.

15

20

When R_1 - R_3 are hydrogen and R_4 is 2-dimethylaminoethoxy, it is preferred that:

- step (i) is performed by reacting the compound of the formula (II) with ethylene carbonate with an alkali metal carbonate (e.g. potassium carbonate) or an alkali earth metal carbonate, as a base and dimethyl sulfoxide as a solvent,
- step (ii) is performed by reacting the compound of the formula (III) with the compound of the formula (IV) in the presence of an alkali metal tert-butoxide (e.g. potassium tert-butoxide) or an alkali earth metal tert-butoxide, in tetrahydrofuran,
- the pharmaceutically acceptable acid of the formula HA in step (iii) is phosphoric acid.

25

30

This invention also features a process for the preparation of compounds of the general formula (I). The process includes the steps of:

- (i) reacting a 3-pyridinol derivative of the general formula (II) or the corresponding hydrochloride with ethylene carbonate, to give another 3-pyridinol derivative of the general formula (III):
- (ii) reacting the 3-pyridinol derivative of the general formula (III) with (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) in the presence of an alkali metal tert-butoxide (e.g., potassium tert-butoxide) or an alkali earth metal tert-butoxide, to give a compound of the general formula (V),
- (iii) converting the compound of the general formula (V) to the compound of the general formula (I) by treatment with a pharmaceutically acceptable acid of the formula HA (e.g., L-malic acid),
- wherein each of R₁, R₂, R₃, and R₄ is hydrogen. Step (i) can be performed in the presence of an alkali metal carbonate (e.g. potassium carbonate) or an alkali earth metal carbonate as a base, and dimethyl formamide as a solvent; and step (ii) can be performed by reacting the compound of the general formula (III) with the compound of the formula (IV) in tetrahydrofuran.

In this process, (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) can be prepared by:

- (iv) acidification of racemic 2-methylpiperazine of the formula (VI) with L-tartaric acid and fractional crystallization to give (*R*)-2-methylpiperazine, L-tartrate of the formula (VII):
- (v) basification of (2*R*)-2-methylpiperazine, L-tartrate of the formula (VII) to give (*R*)-2-methylpiperazine of the formula (VIII):
- (vi) tritylation of (*R*)-2-methylpiperazine of the formula (VIII) to give (*R*)-3-methyl-1-tritylpiperazine of the formula (IX),
- (vii) condensation of (*R*)-3-methyl-1-tritylpiperazine of the formula (IX) with 2,3-dichloropyrazine to give (2*R*)-1-(3-chloro-2-pyrazine)-2-methyl-4-tritylpiperazine of the formula (X):
- (viii) detritylation of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X) to give (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV),
- (ix) and conversion of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) to a suitable acid addition salt (e.g, hydrochloride salt).

The formula (I) to (X) are the same as described above.

This invention further features a process for the preparation of compounds of the general formula (I). The process includes the steps of:

- (i) O-alkylating a 3-pyridinol derivative of the general formula (II) or the corresponding hydrochloride, to give another 3-pyridinol derivative of the general formula (III),
 - (ii) reacting the 3-pyridinol derivative of the general formula (III) with (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) in the presence of potassium tert-butoxide, to give a compound of the general formula (V),
 - (iii) converting the compound of the general formula (V) to the compound of the general formula (I) by treatment with a pharmaceutically acceptable acid of the formula HA (e.g., succinic acid),
- wherein each of R₁, R₂, and R₃ is hydrogen, R₄ is C₁-C₆-alkoxy (e.g., ethoxy). Step (i) can be performed by reacting the hydrochloride of the compound of the general formula (II) with 2-chloroethanol in an aqueous solution of a hydroxide (e.g., sodium hydroxide), and step (ii) can be performed by reacting the compound of the general formula (III) with the compound of the formula (IV) in a solvent system consisting of methyl tert-butylether and tetrahydrofuran.

This invention also features a process for the preparation of compounds of the general formula (I). The process includes the steps of:

- (i) reacting a 3-pyridinol derivative of the general formula (II) or the corresponding hydrochloride with ethylene carbonate, to give another 3-pyridinol derivative of the general formula (III):
- (ii) reacting the 3-pyridinol derivative of the general formula (III) with (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) in the presence of an alkali metal tert-butoxide (e.g., potassium tert-butoxide) or an alkali earth metal tert-butoxide, to give a compound of the general formula (V),
- (iii) converting the compound of the general formula (V) to the compound of the general formula (I) by treatment with a pharmaceutically acceptable acid of the formula HA (e.g., phosphoric acid),

wherein each of R₁, R₂, and R₃ is hydrogen, R₄ is di-C₁-C₆-alkylamino-C₂-C₆-alkoxy (e.g., 2-dimethylaminoethoxy). Step (i) can be performed in the presence of an alkali metal

carbonate (e.g. potassium carbonate) or an alkali earth metal carbonate as a base, and dimethyl sulfoxide as a solvent; and step (ii) can be performed by reacting the compound of the general formula (III) with the compound of the formula (IV) in tetrahydrofuran.

Another object of the present invention is a process for preparing (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine, hydrochloride by reacting (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) with hydrochloric acid.

Another object of the present invention is the compound (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine, hydrochloride.

Another object of the present invention is a compound of the formula (I), wherein R_1 - R_4 are hydrogen and HA is L-malic acid.

Another object of the present invention is a compound of the formula (I), wherein R_1 - R_3 are hydrogen, R_4 is ethoxy and HA is succinic acid.

Another object of the present invention is a compound of the formula (I), wherein R_1 - R_3 are hydrogen, R_4 is 2-dimethylaminoethoxy, and HA is phosphoric acid.

Also within the scope of this invention is a method for the treatment or prophylaxis of a serotonin-related disorder, particularly 5-HT_{2C} receptor related. The method includes administering to a subject in need thereof an effective amount of a compound of the formula (I), wherein R_1 - R_3 are hydrogen, R_4 is selected from hydrogen, ethoxy, and 2-dimethylaminoethoxy, and HA is selected from L-malic acid, succinic acid, and phosphoric acid. Examples of such disorders are obesity and type II diabetes.

Also within the scope of this invention is the use of a compound of formula (I), wherein R_1 - R_3 are hydrogen, R_4 is selected from hydrogen, ethoxy, and 2-dimethylaminoethoxy, and HA is selected from L-malic acid, succinic acid, and phosphoric acid, to manufacture a medicament for the treatment or prophylaxis of a serotonin-related disorder, particularly 5-HT_{2C} receptor related. Examples of such disorders are obesity and type II diabetes.

Detailed description of the invention

Below, the various terms used in the above definition of the compounds of the compounds of the general formulas (I)-(V) will be explained.

C₁-C₆-alkyl is a straight or branched alkyl group having 1-6 carbon atoms. Exemplary alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, n-hexyl, and isohexyl.

For parts of the range "C₁-C₆-alkyl" all subgroups thereof are contemplated such as C₁-C₅-alkyl, C₁-C₄-alkyl, C₁-C₃-alkyl, C₁-C₂-alkyl, C₂-C₆-alkyl, C₂-C₅-alkyl, C₂-C₄-alkyl, C₂-C₃-alkyl, C₃-C₆-alkyl, C₄-C₅-alkyl, etc.

C₁-C₆-alkoxy is a straight or branched alkyl group having 1-6 carbon atoms.

5 Exemplary alkoxy groups include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, n-pentoxo, isopentoxo, n-hexoxo, and isohexoxo.

For parts of the range "C₁-C₆-alkoxy" all subgroups thereof are contemplated such as C₁-C₅-alkoxy, C₁-C₄-alkoxy, C₁-C₃-alkoxy, C₁-C₂-alkoxy, C₂-C₆-alkoxy, C₂-C₅-alkoxy, C₂-C₄-alkoxy, C₂-C₃-alkoxy, C₃-C₆-alkoxy, C₄-C₅-alkoxy, etc.

10 The term "DSC" in the present description means "differential scanning calorimetry".

The term "DVS" in the present description means "dynamic vapor sorption gravimetry".

The term "FBE" in the present description means "free base equivalents".

15 The term "halogen" in the present description is intended to include fluorine, chlorine, bromine and iodine.

The term "IDR" in the present description means "intrinsic dissolution rate".

The term "IP" in the present description means "in process".

The term "NLT" in the present description means "not less than".

20 "Pharmaceutically acceptable acid salts" mean salts which are pharmaceutically acceptable, as defined above, and which possess the desired pharmacological activity. Such salts include acid addition salts formed with any organic and inorganic pharmaceutically acceptable acid, such as hydrogen chloride, hydrogen bromide, hydrogen iodide, sulfuric acid, phosphoric acid, acetic acid, glycolic acid, maleic acid, malonic acid, oxalic acid, toluenesulfonic acid, methanesulfonic acid, trifluoroacetic acid, fumaric acid, succinic acid, 25 malic acid, tartaric acid, citric acid, benzoic acid, ascorbic acid and the like.

DMF means dimethyl formamide, DMSO means dimethyl sulfoxide, IPA means isopropanol, MtBE means methyl tert-butyl ether, RH means relative humidity, RT means room temperature, t-BuOK means tert-butyl alcohol, potassium salt, THF means 30 tetrahydrofuran, and trityl means triphenylmethyl.

It should be noted that both E- and Z-isomers of the compounds, optical isomers, as well as mixtures thereof, and all isotopes are included within the scope of the invention. By the expression "isotopes" is meant all compounds with naturally occurring isotopes such as all possible deuterium and ¹³C-isotopes of the compounds according to the invention. The

compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constituted such compounds. For example, the compounds may be radiolabelled with radioactive isotopes, such as for example tritium (^3H), ^{125}I or ^{14}C . All isotopic variations of the compounds of the present invention, whether
5 radioactive or not, are intended to be encompassed within the scope of the present invention.

The chemicals used in the above-described synthetic routes may include, for example, solvents, reagents, catalysts, protecting group and deprotecting group agents. The methods described above may also additionally include steps, either before or after the steps described specifically herein, to add or remove suitable protecting groups in order to
10 ultimately allow synthesis of the compounds of the formula (I). Synthetic chemistry transformations and protecting group methodologies (protection and deprotection) useful in synthesizing applicable compounds of the formula (I) are known in the art and include, for example, those described in R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*,
15 2nd Ed., John Wiley and Sons (1991); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); and L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995) and subsequent editions thereof.

In accordance with the present invention, the compound of the formula (I), wherein $\text{R}_1\text{-R}_3$ are hydrogen, R_4 is selected from hydrogen, ethoxy or 2-dimethylaminoethoxy and HA
20 is selected from L-malic acid, succinic acid, and phosphoric acid, can be brought into suitable galenic forms, such as compositions for oral use, for injection, for nasal spray administration or the like, in accordance with accepted pharmaceutical procedures. Such pharmaceutical compositions according to the invention comprise an effective amount of the aforementioned compound in association with compatible pharmaceutically acceptable carrier materials, or
25 diluents, as are well known in the art. The carriers may be any inert material, organic or inorganic, suitable for enteral, percutaneous, subcutaneous or parenteral administration, such as: water, gelatin, gum arabicum, lactose, microcrystalline cellulose, starch, sodium starch glycolate, calcium hydrogen phosphate, magnesium stearate, talcum, colloidal silicon dioxide, and the like. Such compositions may also contain other pharmacologically active agents, and
30 conventional additives, such as stabilizers, wetting agents, emulsifiers, flavouring agents, buffers, and the like.

The compositions according to the invention can e.g. be made up in solid or liquid form for oral administration, such as tablets, pills, capsules, powders, syrups, elixirs, dispersible granules, cachets, suppositories and the like, in the form of sterile solutions.

suspensions or emulsions for parenteral administration, sprays, e.g. a nasal spray, transdermal preparations, e.g. patches, and the like.

As mentioned above, the aforementioned compounds of the invention may be used for the treatment or prophylaxis of a subject (e.g., a human or an animal) suffering from a serotonin-related disorder or conditions, particularly 5-HT_{2C} receptor related, such as memory disorders, such as Alzheimer's disease; schizophrenia; mood disorders such as depression; anxiety disorders; pain; substance abuse; sexual dysfunctions such as erectile dysfunction; epilepsy; glaucoma; urinary disorders, such as urinary incontinence; menopausal and post-menopausal hot flushes; eating disorders, such as binge eating disorders, anorexia nervosa and bulimia; weight gain associated with antipsychotic drug administration, type II diabetes; and particularly obesity.

The method of treatment or prophylaxis includes administering to a subject in need of treatment an effective amount of the compound of this invention. The term "treating" or "treated" refers to administering a compound of this invention to a subject with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect a disease, the symptoms of the disease or the predisposition toward the disease. "An effective amount" refers to an amount of a compound which confers a therapeutic effect on the treated subject. The therapeutic effect may be objective (i.e., measurable by some test or marker) or subjective (i.e., subject gives an indication of or feels an effect). The dose level and frequency of dosage of the specific compound will vary depending on a variety of factors including the potency of the specific compound employed, the metabolic stability and length of action of that compound, the patient's age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the condition to be treated, and the patient undergoing therapy. The daily dosage may, for example, range from about 0.001 mg to about 100 mg per kilo of body weight, administered singly or multiply in doses, e.g. from about 0.01 mg to about 25 mg each. Normally, such a dosage is given orally but parenteral administration may also be chosen.

The invention also refers to the use of the aforementioned compounds to manufacture a medicament for the treatment or prophylaxis of a serotonin-related disorder, particularly 5-HT_{2C} receptor related.

All publications mentioned herein are hereby incorporated by reference. By the expression "comprising" we understand including but not limited to.

The invention will now be illuminated by the following Examples, which are only intended to illuminate and not restrict the invention in any way. Examples 1A-3A illustrate

steps (i)-(iii) when R₁-R₄ are hydrogen. Examples 1B-3B illustrate steps (i)-(iii) when R₁-R₃ are hydrogen and R₄ is ethoxy. Examples 1C-3C illustrate steps (i)-(iii) when R₁-R₃ are hydrogen and R₄ is 2-dimethylaminoethoxy. Examples 4-8 illustrate steps (iv)-(viii) when R₁-R₃ are hydrogen and R₄ is selected from the group consisting of hydrogen, ethoxy, and 2-dimethylaminoethoxy. Regarding the compound of the formula (IV) as prepared in step (viii), it has been shown that this compound is not stable for storage as a free base. However, the corresponding hydrochloride thereof is considerably more stable. Example 9 describes the preparation of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine, hydrochloride. Furthermore, the properties of different salts of the formula (I) were evaluated. Example 10 describes the preparation of (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, fumarate. By making the fumarate in Example 10, a more pure product is obtained before converting to the L-malate salt (prepared in Example 3A). Example 11 describes the comparison of the properties of different salts of Example 2A. Example 12 describes the comparison of the properties of different salts of Example 2B. Example 13 describes the comparison of the properties of different salts of Example 2C.

EXAMPLES

General

2-ethoxy-3-hydroxypyridine, HCl (for Example 1B) and 2-(2-dimethylaminoethoxy)pyridin-3-ol (for Example 1C) were purchased from Nordic Synthesis in Karlskoga, Sweden. The other chemicals mentioned are commercially available and could e.g. be purchased from Aldrich. For Examples 1B-3B, HPLC analysis confirmed that the desired products were obtained. The HPLC pump was a Varian 9012. A Varian 9050 detector at 220 nm was used. The eluent was 80 mM KH₂PO₄ in a 80:20 mixture of water/acetonitrile. The column was a Varian C8, 150 x 4.5 mm. For Examples, 1C-3C, HPLC analysis also confirmed that the desired products were obtained. The eluent was A: 0.1% trifluoroacetic acid in water, B: acetonitrile; increasing percentage of acetonitrile during 3 minutes. The column was a YMC FL-ODS AQ S-5 μ , 12 nm, 50x4.6mm (Example 1C) and a Hypersil BDS C18, 3 μ , 30x4.6mm (Example 2C-3C), respectively.

EXAMPLE 1A – Preparation of 2-(3-pyridinyloxy)ethanol

3-hydroxypyridine (20.0 kg), ethylene carbonate (19.4 kg), K₂CO₃ (18.9 kg), and DMF (75.5 kg) were charged to a 400 L reactor and heated to 86°C. After 13 h, an aliquot was

taken. There was still some 3-hydroxypyridine left by GC so more ethylene carbonate (1.0 kg) was added. After 20 h, the reaction was deemed complete. The reaction was cooled to 20°C, H₂O (85 L) was slowly added to the 400 L reactor, and stirring continued for 30 min. The title compound was extracted with 35°C CH₂Cl₂ (4 x 170 L) sending the CH₂Cl₂ extractions into
5 the 1000 L portable tank. The aqueous layer was discarded, and the 400 L reactor was cleaned. The solution of the title compound in the 1000 L portable tank was transferred to the 400 L and distilled in the 400 L reactor sending the distillate to the 400 L receiver. Toluene (200 L) was added to the 400 L reactor and distilled sending the distillate to the 400 L receiver. The resulting oil was transferred to a 20 L glass bottle to produce 21.0 kg (72%) of
10 the title compound of 100.0% GC purity.

The yield of the title compound according to D1, starting from 3-hydroxypyridine, 2-chloroethanol and K₂CO₃ in DMF was 19%. The title compound has also been prepared according to EP 0 286 242 A2 starting from methyl [(3-pyridinyl)oxy]acetate.

15 ¹H NMR (400 MHz, CDCl₃): δ 8.30 (1H, br s), 8.20 (1H, t, J = 3.1 Hz), 7.23 (2H, t, J = 2.1 Hz), 4.13 (2H, t, J = 4.6 Hz), 3.99 (2H, t, J = 4.6 Hz), 3.76 (1H, br s), 0.00 (TMS, reference).
¹³C NMR (100 MHz, CDCl₃): δ 155.09 (s), 142.09 (d), 137.95 (d), 123.99 (d), 121.32 (d), 69.83 (t), 60.98 (t), 0.00 (TMS, reference).
IR (liq.) 2382 (w), 2082 (w), 1996 (w), 1954 (w), 1587, 1576 (s), 1488, 1478 (s), 1429 (s),
20 1272 (s), 1235 (s), 1084, 1054 (s), 802, 707 (s), cm⁻¹.
Anal. Calcd for C₇H₉NO₂: C, 60.42; H, 6.52; N, 10.07. Found: C, 60.14; H, 6.51; N, 10.39.
HRMS (FAB) calcd for C₇H₉NO₂ +H₁ 140.0712, found 140.0705.

EXAMPLE 1B – Preparation of 2-[(2-ethoxy-3-pyridinyl)oxy]ethanol

25 To water (30.0 kg) was added 3.40 kg of NaOH. The resulting suspension was stirred until complete dissolution had occurred. The temperature of the solution was then adjusted to approx. 60°C. To the alkaline solution was carefully added 2-ethoxy-3-hydroxypyridine, HCl (6.00 kg). The temperature was raised to 85.0°C and the mixture stirred for 30 min to allow for complete dissolution. During 85 min, 2-chloroethanol (4.10 kg) was added, while the
30 temperature was kept between 88.0-92.0°C. The reaction was allowed to age for 50 min at approx. 90°C, after which time HPLC indicated complete conversion of 2-ethoxy-3-hydroxypyridine, HCl to the title compound. The reaction mixture was poured into 65.1 kg of methyl tert-butylether (MtBE). The reaction vessel was rinsed with water (2.0 kg). To the resulting two-phase system was added 3.05 kg of NaCl and 0.20 kg of NaOH. The resulting

mixture was stirred for 55 min, while the temperature was lowered to 24.9°C. The mixture was allowed to stir for another 20 min at 23.7-24.9°C. The water phase was removed. The remaining organic phase was concentrated by distillation at atmospheric pressure. When 45L had been removed, the residue was cooled to room temperature and transferred to a drum. The residue weighed 33.1 kg. According to HPLC, the assay is 118.9 g of the title compound per kg of mixture, corresponding to a yield of 6.25 kg (quantitative yield) of pure title compound.

The yield of the title compound according to D1, starting from 2-bromo-3-(2-[[tert-butyl(dimethyl)silyl]oxy]ethoxy)pyridine and sodium ethoxide in ethanol was 41%.

10 EXAMPLE 1C – Preparation of 2-{[2-(2-dimethylaminoethoxy)-3-pyridinyl]oxy}ethanol

DMSO (17L) and potassium carbonate (8.2kg) were charged to the 100L-glasslined reactor. The mixture was heated to 58°C while stirring. 2-(2-Dimethylaminoethoxy)pyridin-3-ol (4.11kg) was charged to the reactor during 25 minutes (gas evolution). The mixture was heated to 117°C before the prepared ethylene carbonate solution was added through the manhood in portions. After 1L ethylene carbonate solution was added IP-HPLC was taken to ensure that the title compound had started to form. The total addition of ethylene carbonate solution was made during 1h 35 minutes. Temperature after the addition was 124°C. A HPLC sample was taken from the reaction mixture 10 minutes after finished addition. It showed 95% conversion to product. The reaction-mixture was allowed to cool to 70°C before water (45L) and sodium chloride (4.06kg) was charged to the reactor. The mixture was then stirred for 20 minutes at 70°C before cooled to RT. Ethyl acetate (137L) was charged into a 328L glass-lined reactor. The water mixture was transferred from the 100L reactor to the 328L reactor. The mixture was stirred for 25 minutes before the mechanical stirrer was turned off. After allowing the phases to separate for 40 minutes, the water-phase (65L) was discharged. The mixture was heated to reflux and 102L of ethyl acetate was distilled off. Toluene (72L) was added and the mixture was heated to reflux again and 61L of toluene/ethyl acetate was distilled off. The reaction mixture was then cooled to RT. The title compound was never isolated; it was directly used in Example 2C.

30 The preparation of the title compound has not been described in D1.

EXAMPLE 2A – Preparation of (2R)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine

A 400 L reactor containing a toluene solution of (2R)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine (for preparation of this compound, see Example 8 below) was distilled to
5 remove toluene. THF (40 kg) was added, and the slurry was stirred until all the oil dissolved. The solution was transferred to a drum with a THF rinse (10 kg). The THF/toluene solution of (2R)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine (ca 9.7 kg thereof in 90.4 kg solution) was transferred to a 1200 L reactor with a THF rinse (4 L). 2-(3-pyridinyloxy)ethanol from
10 Example 1A (7.6 kg) was transferred to the 1200 L reactor with a THF rinse (34 L) and stirred at room temperature for 15 min. 20wt% KOt-Bu (33.3 kg) was added followed by a THF rinse (4 kg). The reaction was monitored by GC and was complete in 15 min. The reaction in the 1200 L reactor was quenched with H₂O (119 L) and extracted with CH₂Cl₂ (1 x 215 L, 2 x 117 L) sending the CH₂Cl₂ extractions into the 800 L receiver. The CH₂Cl₂ solution was transferred to the 1200 L reactor and the CH₂Cl₂ was removed by distillation. A
15 solvent swap distillation using MeOH (2 x 215 L) was performed distilling to a volume of 100 L. The MeOH solution of the title compound was filtered through a bag filter containing carbon (5 kg) recycling the solution four times from the 1200 L reactor to the 800 L receiver. The solution in the 800 L receiver was transferred to the 1200 L reactor with a MeOH rinse (50 L). The solution was distilled to a final volume of 70 L and transferred to a drum with a
20 MeOH rinse (10 L) in preparation for the compound in Example 3A. The yield was quantitative.

The yield of the hydrochloride of the title compound according to D1, starting from 3-chloro-2-[4-tert-butoxycarbonyl-(2R)-methyl-1-piperazinyl]pyrazine and 2-(3-pyridinyloxy)ethanol with potassium tert-butoxide in dioxane was 31%.

25

¹H NMR (400 MHz, CDCl₃): δ 8.35 (1H, br s), 8.25 (1H, t, J = 3.0 Hz), 7.74 (1H, d, J = 3.0 Hz), 7.49 (1H, d, J = 2.5 Hz), 7.23 (2H, br t, J = 2.3 Hz), 4.76 (1H, dt, J = 12.2, 4.8 Hz), 4.67 (1H, dt, J = 11.7 Hz, 5.2 Hz), 4.49-4.43 (1H, m), 4.40 (2H, t, J = 4.8 Hz), 3.72 (1H, br d, J = 12.8 Hz), 3.27 (1H, td, J = 11.7, 3.0 Hz), 3.06 (1H, dd, J = 12.3, 3.6 Hz), 2.99 (1H, br d, J =
30 12.2 Hz), 2.88 (1H, td, J = 11.7, 3.4 Hz), 2.75 (1H, br d, J = 12.3 Hz), 1.72 (1H, br s), 1.19 (3H, d, J = 6.6 Hz), 0.00 (TMS, reference).

¹³C NMR (100 MHz, CDCl₃): δ 154.86 (s), 150.35 (s), 147.30 (s), 142.60 (d), 138.09 (d), 134.14 (d), 129.89 (d), 123.87 (d), 121.17 (d), 66.54 (t), 63.97 (t), 50.86 (t), 49.52 (d), 46.22 (t), 42.78 (t), 14.26 (q), 0.00 (TMS, reference).

IR (liq.) 2068 (w), 1996 (w), 1576 (s), 1528 (s), 1475 (s), 1430 (s), 1330, 1276 (s), 1268 (s), 1246, 1208 (s), 1187 (s), 1181 (s), 1063, 708, cm^{-1} .

HRMS (FAB) calcd for $\text{C}_{16}\text{H}_{21}\text{N}_5\text{O}_2 + \text{H}_1$ 316.1773, found 316.1777.

$[\alpha]_D^{25} = -24^\circ$ (c 0.92, water).

- 5 Anal. Calcd for $\text{C}_{16}\text{H}_{21}\text{N}_5\text{O}_2$: C, 60.94; H, 6.71; N, 22.21. Found: C, 59.98; H, 6.73; N, 22.40.

EXAMPLE 2B – Preparation of (2R)-1-(3-{2-[(2-ethoxy-3-pyridinyl)oxy]ethoxy}-2-pyrazinyl)-2-methylpiperazine

- (2R)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine (9.00 kg) (see Example 8 for
10 preparation) was dissolved in 20.1 kg of water by stirring at room temperature for 25 min. Activated charcoal (0.50 kg) and celite (0.50 kg) was added. The resulting suspension was stirred for 30 min at room temperature. The charcoal and celite was removed by filtration under pressure. The filtered water-phase was poured into 63.9 kg of MtBE (methyl tert-butylether). To the resulting two-phase system were added NaOH (1.50 kg) and NaCl (4.40
15 kg). After stirring for 25 min at room temperature, the water phase was discarded. The remaining organic phase was concentrated by distillation at atmospheric pressure. A total of 45L was removed. To the remaining content of the reactor was added an MtBE-solution of 2-[(2-ethoxy-3-pyridinyl)oxy]ethanol (one complete batch from Example 1B). The resulting mixture is concentrated by distillation at atmospheric pressure. A total of 40L was removed.
20 To the residue, 40.8 kg of MtBE was added. Further concentration by distillation was done with a total of 20L being removed. A solution of potassium tert-butoxide (KtBuO) was made by suspending 2.96 kg of KtBuO in 13.1 kg of THF. After stirring for 30 min at room temperature almost everything was in solution. After emptying the reactor, a second solution was made by in a similar way (3.02 kg of KtBuO to 13.0 kg of THF). The mixture of 2-[(2-ethoxy-3-pyridinyl)oxy]ethanol and (2R)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine was
25 heated to 51.0°C. To this mixture was added the solutions of KtBuO in THF. During the addition, the temperature was kept between 50.7-55.6°C. The total addition time was 50 min. After aging for 35 min at 55°C, HPLC indicated that the reaction was complete. The reaction mixture was cooled to room temperature and 23.1 kg of water was cautiously added while
30 maintaining the temperature below 25°C. After stirring for 15 min at approx. 23°C, the water phase was discarded. The remaining organic phase was concentrated by distillation at atmospheric pressure. After 45L had been removed, heptane (38.0 kg) was added. Further distillation removed 50L, followed by addition of heptane (33.8 kg). By removing another 30L via distillation, all remaining MtBE had been removed from the heptane solution. To the

residue, 20.8 kg of water was added and the temperature raised to 72.8°C. After stirring for 15 min, the water phase was removed at this temperature. The remaining solution was concentrated by distillation at atmospheric pressure. A total of 50L was removed. The residue was cooled to 50.5°C. The solution was seeded with 5.0 g of the title compound to induce crystallization. The resulting suspension was stirred at 51.5°C for 35 min while maintaining maximum stirring speed. Cooling was then continued to 8.1°C over night. The product was isolated via filtration under suction. The resulting product cake was washed with 11L of heptane. The semi-dry cake weighing 9.63 kg was dried at 40°C/vakuum. The dried product weighed 8.1 kg (66%).

The yield of the fumarate of the title compound according to D1, starting from (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine and 2-[(2-ethoxy-3-pyridinyl)oxy]ethanol with sodium *tert*-butoxide in dioxane was 17%.

EXAMPLE 2C – Preparation of (2*R*)-1-[3-(2-{[2-(2-dimethylaminoethoxy)-3-pyridinyl]oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine

(2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine (20L in toluene solution; see Example 8 for preparation) and active charcoal (201g) was added to the 328L-reactor containing 2-[(2-(2-dimethylaminoethoxy)-3-pyridinyl)oxy]ethanol (Example 1C) in toluene solution. The mixture was stirred for 25 minutes before transferred to a through a GAF-filter (containing a Teflon filter bag) to a 250L-stainless steel reactor. The mixture was heated to 100°C before the prepared “THF/ potassium *tert*-butoxide –solution” was added to the reactor using a 50L addition bowl. After 10L “THF/ potassium *tert*-butoxide –solution” was added, an IP-HPLC was taken to ensure that the title compound had started to form. The total addition of THF/ potassium *tert*-butoxide -solution was made during 2h and 15 minutes.

Water (21L) and sodium chloride (1kg) was added to a 198L-glasslined reactor. The salt was dissolved before the reaction mixture was transferred from the 328L-reactor and quenched into the 198L-reactor. Toluene (10L) was also transferred to make sure that no reaction-mixture was left in the hoses used for the transfer. The mixture was then heated to 70°C before the mechanical stirrer was turned off. The phases were allowed to separate for 30 minutes before the water phase (25L) was discharged. The mixture was heated to reflux and toluene (10L) was distilled off. Water (100L) was added and the mixture was heated to reflux again and 60L of water was distilled off. The remaining solution was then cooled to RT. The title compound was never isolated; it was directly used in the next step (Example 3C).

IP sample (HPLC) of the toluene mixture confirmed the retention time for the desired product ($t_{\text{ret}}=2.2$ minutes).

The preparation of the title compound has not been described in D1.

5

EXAMPLE 3A - Preparation of (2R)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, L-malate salt

To a 120 L receiver was added (2R)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine from Example 2A (4.09 kg) and MeOH (12.7 kg) with stirring for about
10 7 min until all substance had dissolved. The solution was transferred to the 120 L reactor through an in-line filter followed by a MeOH (2 kg) rinse. The L-malic acid (1.79 kg) was added to the 120 L receiver followed by acetone (8.8 kg) with stirring for about 13 min until all the L-malic acid had dissolved. The solution was transferred to the 120 L reactor through an in-line filter followed by an acetone (2 kg) rinse. The solution in the 120 L reactor was
15 cooled to 5°C. MtBE (25 kg) was added to the 120 L receiver and slowly transferred through an in-line filter into the 120 L reactor. The solution was stirred at 0°C for 1 h and filtered onto the 18" nutsche sending the filtrate into the 120 L receiver. (Inspection showed little product in the nutsche; only small amounts of oil were present.) MeOH (40 L) was added to the 18" nutsche to dissolve the oil. The MeOH solution and the filtrate in 120 L receiver were
20 transferred through an in-line filter into 120 L reactor and the solution was distilled to a solid sending the distillate to the 120 L receiver, which was disposed. MeOH (15 kg) was added through an in-line filter to 120 L reactor. The solution in 120 L reactor was distilled atmospherically replacing the MeOH with EtOH (35 kg) to a volume of 15 L. The solution was cooled to 58.6°C, seeded with 10 g of (2R)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, L-malate salt, and cooling continued to 25°C. Stirring was continued
25 for 24 h, and the resulting slurry was filtered onto the 18" nutsche sending the filtrate into the 120 L receiver. The filter cake on the 18" nutsche was rinsed with filtered EtOH (15 kg) sending the rinse into the 120 L receiver. The title compound was dried under vacuum using 60°C nitrogen, deagglomerated, and packaged to yield 3.58 kg (61%) of the title compound of
30 98.61% GC purity.

The preparation of the title compound has not been described in D1.

M.Pt. 124.5-126.6°C.

¹H NMR (400 MHz, d₄-MeOH): δ 8.28 (1H, d, J = 2.6 Hz), 8.16 (1H, d, J = 4.6 Hz), 7.78 (1H, d, J = 2.5 Hz), 7.65 (1H, d, J = 3.0 Hz), 7.49 (1H, dd, J = 8.6, 2.5 Hz), 7.38 (1H, dd, J = 8.3, 4.8 Hz), 4.78-4.73 (2H, m), 4.65 (1H, quintet, J = 3.4 Hz), 4.48 (2H, t, J = 4.4 Hz), 4.29 (1H, dd, J = 7.3, 5.3 Hz), 3.98 (1H, dt, J = 14.3, 3.1 Hz), 3.50 (1H, ddd, J = 14.3, 11.2, 2.6 Hz), 3.31 (d₄-MeOH), 3.30-3.24 (2H, m), 3.21-3.15 (2H, m), 2.77 (1H, dd, J = 15.8, 5.1 Hz), 2.53 (1H, dd, J = 15.8, 7.1 Hz), 1.28 (3H, d, J = 7.1 Hz), 0.00 (TMS, reference).

¹³C NMR (100 MHz, d₄-MeOH): δ 179.64 (s), 176.38 (s), 157.02 (s), 152.32 (s), 147.25 (s), 142.72 (d), 138.58 (d), 135.14 (d), 133.35 (d), 125.97 (d), 123.40 (d), 69.67 (d), 68.00 (t), 65.78 (t), 48.61 (d), 48.32 (t), 44.31 (t), 41.84 (t), 40.20 (t), 14.80 (q), 0.00 (TMS, reference).

IR (diffuse reflectance) 3030 (s), 3009, 2970, 2469 (b), 2358, 2342 (b), 2318 (b), 2303, 1571 (s), 1471, 1449, 1408, 1275 (s), 1233 (s), 1206 (s), cm⁻¹.

HRMS (FAB) calcd for C₁₆H₂₁N₅O₂ + H₁ 316.1773, found 316.1780.

[α]_D²⁵ = -21° (c 0.98, water).

Anal. Calcd for C₁₆H₂₁N₅O₂·C₄H₆O₅: C, 53.44; H, 6.05; N, 15.58. Found: C, 53.23; H, 6.08; N, 15.29.

EXAMPLE 3B – Preparation of (2R)-1-(3-{2-[(2-ethoxy-3-pyridinyl)oxy]ethoxy}-2-pyrazinyl)-2-methylpiperazine, succinate salt

(2R)-1-(3-{2-[(2-ethoxy-3-pyridinyl)oxy]ethoxy}-2-pyrazinyl)-2-methylpiperazine (8.00 kg) obtained in Example 2B was dissolved in 24.0 kg of 1-propanol by heating to 60.5°C. The undissolved material was removed by filtration. Succinic acid (1.30 kg) was dissolved in 23.9 kg of 1-propanol by heating to 60.6°C. All undissolved material was removed by filtration. The procedure was repeated once with another 1.30 kg of succinic acid and 23.9 kg of 1-propanol. All of the above solutions were mixed in a reactor. After mixing, the content's temperature was 53.9°C. The solution was heated to 78.3°C. Cooling was applied. At 40.3°C, seeding with the title compound was done. Crystallization started after the seeding. The contents were stirred, while the temperature was lowered to 18.3°C during the night. The product was isolated via filtration under suction. The resulting product cake was washed with 10L of 1-propanol. The resulting semi-dry cake, weighing 17.2 kg was dried at 50°C/vakuum. After drying, the product, the title compound weighed 9.16 kg (86%).

The preparation of the title compound has not been described in D1.

Melting temperature onset, 123°C; thermal melting range, 116-132°C.

¹H NMR (500 MHz, DMSO-d₆): δ 1.12 (d, J = 6.8 Hz), 1.28 (t, J = 7.1 Hz), 2.34 (s), 2.79 (m), 2.79 (m), 2.95 (m), 2.95 (m), 3.20 (dddd, J = 13.7, 11.5, 3.0, 1.1 Hz), 3.77 (dt, J = 3.0, 13.7 Hz), 4.30 (q, J = 7.1 Hz), 4.37 (strong coupled), 4.49 (dq, J = 7.0, 6.8, 3.1 Hz), 4.62 (strong coupled), 6.89 (dd, J = 5.0, 7.8 Hz), 7.33 (dd, J = 1.5, 7.8 Hz), 7.58 (d, J = 2.8 Hz), 7.69 (dd, J = 1.5, 5.0 Hz), 7.77 (d, J = 2.8 Hz).

¹³C-NMR (500 MHz, DMSO-d₆): δ 14.2, 14.5, 30.5, 40.5, 44.0, 47.9, 48.4, 61.0, 64.0, 66.5, 116.8, 119.9, 130.3, 133.7, 137.2, 142.7, 146.3, 149.9, 153.8, 174.5.

Anal. Calcd for C₁₈H₂₅N₅O₃ · C₄H₆O₄: C, 55.34%; H, 6.54%; N, 14.67%. Found: C, 54.86%; H, 6.75%; N, 13.85%.

EXAMPLE 3C – Preparation of (2R)-1-[3-(2-{[2-(2-dimethylaminoethoxy)-3-pyridinyl]oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, phosphoric acid salt

(2R)-1-[3-(2-{[2-(2-dimethylaminoethoxy)-3-pyridinyl]oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine (Example 2C) in water solution was added to the 198L-glasslined reactor and heated to reflux. Water (30L) was removed by distillation. The mixture was cooled to RT. before active charcoal (403.5g) was added. The mixture was stirred for 30 minutes before transferred to the 100L glass-lined reactor using Teflon hose and a GAF-filter (containing a polypropylene filter bag) and a Millipore filter (containing a polyguard filter cartridge). The hoses and filters were rinsed with water (10L). To the water solution in the 100L reactor phosphoric acid (3.83kg) was added. The mixture was heated to reflux and water was distilled off until an 18L solution of (2R)-1-[3-(2-{[2-(2-dimethylaminoethoxy)-3-pyridinyl]oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine in water/phosphoric acid remained. Ethanol (71L) was added to the solution while heating the mixture to reflux. All (2R)-1-[3-(2-{[2-(2-dimethylaminoethoxy)-3-pyridinyl]oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine was in solution before the heat was turned off and the mixture allowed to cool to RT. A small amount of the solution was taken out to create seeds and the reaction-mixture was seeded when the temperature was 60-68°C. The mixture was allowed to crystallize for 3 days (over a weekend, no cooling media was used) and the crystals were collected by filtration using a Teflon filter nutch. The crystals were then dried in a vacuum oven for 2 days at 60°C and at 100-200mbar. 4.5kg of the title compound was isolated. IP sample (HPLC) of the title compound confirmed the retention time for the title compound (t_{ret}=2.2 minutes). The yield of the title compound was 57% starting from 2-(2-dimethylaminoethoxy)pyridin-3-ol.

By starting from (2*R*)-1-[3-(2-[[2-(2-dimethylaminoethoxy)-3-pyridinyl]oxy]-ethoxy)-2-pyrazinyl]-2-methylpiperazine (Example 2C) with a purity of 60-70%, the title compound was obtained with a purity of 99% in one crystallization step.

- 5 The preparation of the title compound has not been described in D1.

Melting temperature onset, 148.0°C (extrapolated); thermal melting range, 140-168°C.

¹H NMR (500 MHz, DMSO-*d*₆): δ 1.21 (d, *J* = 7.0), 2.60 (s), 2.92 (td, *J* = 12.2, 3.2 Hz), 3.02 (dd, *J* = 13.0, 2.6 Hz), 3.08 (dd, *J* = 13.0, 4.2 Hz), 3.16 (m), 3.17 (t, *J* = 5.7 Hz), 3.38 (ddd, *J* = 14.2, 11.6, 2.8 Hz), 3.94 (dt, *J* = 14.2, 3.2 Hz), 4.40 (m), 4.52 (m), 4.59 (m), 4.61 (m), 4.67 (ddd, *J* = 12.3, 5.1, 3.5 Hz), 6.96 (dd, *J* = 7.8, 5.0 Hz), 7.39 (dd, *J* = 7.8, 1.4 Hz), 7.62 (d, *J* = 2.9 Hz), 7.71 (dd, *J* = 5.0, 1.4 Hz), 7.80 (d, *J* = 2.9 Hz).

¹³C-NMR (400 MHz, DMSO-*d*₆): δ 14.3, 38.6, 42.2, 43.4, 46.2, 46.7, 55.5, 61.3, 64.0, 66.4, 117.7, 120.0, 130.9, 133.8, 137.0, 142.7, 145.7, 150.0, 152.8.

15 HRMS *m/z* calcd for C₂₀H₃₀N₆O₃ (M)⁺ 402.2379, found 402.2366.

Anal. Calcd for C₂₀H₃₀N₆O₃ · 3H₃PO₄: C, 34.49%; H, 5.64%; N, 12.07%; O, 34.46%; P, 13.34%. Found: C, 37.20%; H, 6.0%; N, 12.5%; O, 32.4%; P, 12.1%.

EXAMPLE 4 – Preparation of (R)-2-methylpiperazine, L-tartrate

20 To the 1200 L reactor was added 60°C racemic 2-methylpiperazine (100 kg) from a drum. H₂O (240 L) was added, and the solution was cooled to 13°C. To the 1200 L receiver was added L-tartaric acid (150 kg). H₂O (140 L) was added, and the slurry was stirred for 1 h 35 min until dissolution of the solids was complete. The L-tartaric acid solution was transferred to the 1200 L reactor over 2 h while maintaining a temperature of 10-22°C in the
25 1200 L reactor followed by a H₂O rinse (20 L). Ethanol (163 kg) was added to the 1200 L reactor, and the solution was cooled to 2°C. The resulting slurry was stirred for 2 h at 2°C, and filtered through a 36" Nutsche filter sending the filtrate into the 1200 L receiver. The 1200 L reactor and 36" Nutsche filter were washed with H₂O (200 L), and the solids were dried to yield 214 kg of 12% ee (171% based on the title compound). These solids were recharged to a
30 clean 1200 L receiver and H₂O (630 L) was added to the 1200 L receiver, which was heated to 85°C until all the solids had dissolved. The solution was filtered through an in-line filter (C) into the 1200 L reactor, cooled to 5°C, and stirred for 2 h. The resulting slurry was filtered through a clean 36" Nutsche filter sending the filtrate into the 1200 L receiver. The 1200 L reactor and 36" Nutsche filter were washed with H₂O (200 L), and the solids were dried to

yield 104 kg of 93% ee (83% based on the title compound). These solids were recharged to a clean 1200 L receiver and H₂O (254 L) was added to the 1200 L receiver, which was heated to 85°C until all the solids had dissolved. The solution was filtered through an in-line filter (C) into the 1200 L reactor, cooled to 5°C, and stirred for 2 h. The resulting slurry was filtered
5 through a clean 36" Nutsche filter sending the filtrate into the 1200 L receiver. The 1200 L reactor and 36" Nutsche filter were washed with H₂O (200 L), and the solids were dried to yield 92 kg of 99% ee (74% based on the title compound).

D1 does not describe the preparation of the title compound but makes reference to *J. Med. Chem.* 1990, 33, 1645-1656 (D2). The yield of the title compound according to D2,
10 starting from racemic 2-methylpiperazine was 35%.

M.Pt. 255.0-257.0°C.

¹H NMR (400 MHz, D₂O): δ 4.79 (D₂O, reference), 4.36 (2H, s), 3.73-3.64 (4H, m), 3.43 (1H, td J = 13.7, 3.0 Hz), 3.34 (1H, td, J = 12.7, 3.1 Hz), 3.17 (1H, dd, J = 14.2 12.8 Hz), 1.41
15 (3H, d, J = 6.1 Hz), 0.00 (TMS, reference).

¹³C NMR (100 MHz, D₂O): δ 178.46 (s), 73.91 (d), 49.02 (d), 49.00 (MeOH, reference), 45.82 (t), 40.56 (t), 40.10 (t), 15.42 (q).

IR (diffuse reflectance) 3426 (s), 3011 (s), 2999 (s), 2888 (s), 2785 (s,b), 2740 (s,b), 2703 (s,b), 2649 (s,b), 2483 (s,b), 2483 (s,b), 2361 (s), 2354, 2340, 2248, 1638 (s),
20 cm⁻¹.

HRMS (FAB) calcd for C₅H₁₂N₂ + H₁ 101.1079, found 101.1080.

[α]_D²⁵ = 24° (c 1.00, water).

Anal. Calcd for C₄H₆O₆.C₅H₁₂N₂: C, 43.20; H, 7.25; N, 11.19. Found: C, 41.25; H, 7.45; N, 10.71.

25 EXAMPLE 5 – Preparation of (R)-2-methylpiperazine

(R)-2-methylpiperazine, L-tartrate from Example 4 in H₂O (182 L), and Branched octanes (200 L) were added to a 4000 L reactor and stirred until dissolved. More branched octanes (530 L) were added to the 4000 L reactor followed by 50% NaOH (1120 kg) at a
30 temperature between 35°C and 52°C. The solution was heated to 80°C and stirred for 2h. The solution was settled and the lower aqueous phase was transferred to the 4000 L receiver. The solution in the 4000 L reactor was cooled to -21°C and filtered onto a 48" Nutsche filter sending the filtrate to a 1200 L reactor. The 4000 L reactor and 48" Nutsche filter were rinsed with branched octanes (300 L). The solids were dried with 25°C nitrogen and collected to

yield 24.9 kg (67%) of the title compound of NLT 99% ee as determined by chiral HPLC assay. The aqueous solution in the 4000 L receiver was adjusted to pH 8.4 with acetic acid (812 kg) before disposal.

The yield of the title compound according to D2, starting from (*R*)-2-methyl-
5 piperazine, L-tartrate was 42%.

M.Pt. 91-93°C.

¹H NMR (400 MHz, CDCl₃): δ 2.97-2.68 (6H, m), 2.35 (1H, dd, J = 11.7, 10.2 Hz), 1.61 (2H, s), 1.00 (3H, d, J = 6.7 Hz), 0.00 (TMS, reference).

10 ¹³C NMR (100 MHz, CDCl₃): δ 54.14 (t), 51.89 (d), 47.43 (t), 46.46 (t), 20.08 (q), 0.00 (TMS, reference). IR (mull) 3220 (s,b), 2819 (s), 2748, 2042 (w), 1995 (w), 1981 (w), 1328, 1279, 1137, 1094, 960, 859 (s), 845 (s), 795 (s), 621 (s), cm⁻¹.

HRMS (FAB) calcd for C₅H₁₂N₂ + H₁ 101.1079, found 101.1080.

[α]_D²⁵ = -17° (c 0.85, CH₂Cl₂).

15 Anal. Calcd for C₅H₁₂N₂: C, 59.96; H, 12.07; N, 27.97. Found: C, 59.25; H, 11.71; N, 27.64.

EXAMPLE 6 – Preparation of (*R*)-3-methyl-1-tritylpiperazine

To the 1200 L reactor was dissolved (*R*)-2-methylpiperazine from Example 5 (25 kg) in CH₃CN (319 kg) at 15°C to 25°C until dissolution was complete (10 min.) After cooling to
20 5°C to 10°C Et₃N (63 kg) was added. To the 1200 L receiver trityl chloride (69.5 kg) was dissolved in CH₂Cl₂ (106 kg) at 15°C to 25°C. The solution in the receiver was transferred to the reactor over 0.5 h with a CH₂Cl₂ rinse (27 kg), and the solution was heated to 20°C to 30°C. The reaction was monitored by GLC and was complete in 1 h. The resulting slurry was cooled to 8°C to 12°C, filtered onto a 48" Nutsche filter, and rinsed with CH₃CN (40 kg at
25 8°C to 12°C). The filter cake was dried using 50°C to 55°C nitrogen to afford 25.26 kg of the by-product Et₃NHCl (74% yield; easy to filter off the by-product). The filtrate was transferred to the 1200 L reactor and cooled further to -8°C to -10°C for 1 h. The resulting slurry was filtered onto a 24" Nutsche filter and rinsed with -8°C to -10°C CH₃CN (24 kg) sending the filtrate and rinse to the 1200 L receiver. The filter cake was dried with 50°C to 55°C nitrogen
30 to afford another 2.98 kg Et₃NHCl (9% yield). The filtrate was transferred to the 1200 L reactor with a CH₃CN (10 kg) rinse, and distilled under vacuum to an oil of the title compound of 97.99% GC purity. The yield was quantitative.

M.Pt. 134-136°C.

¹H NMR (400 MHz, CDCl₃): δ 7.55-7.40 (6H, br s), 7.25 (6H, t, J = 7.9 Hz), 7.14 (3H, t, J = 7.1 Hz), 3.21-3.13 (2H, m), 3.10-2.90 (1H, br s), 2.94 (2H, t, J = 13.0 Hz), 1.60 (1H, br s), 1.48 (1H, br s), 1.15 (1H, br s), 0.94 (3H, d, J = 6.1 Hz), 0.00 (TMS, reference).

¹³C NMR (100 MHz, CDCl₃): δ 129.41 (d), 127.46 (d), 125.96 (d), 125.83 (s), 56.12 (t), 51.23 (d), 48.73 (t), 46.45 (t), 20.05 (q), 0.00 (TMS, reference).

IR (diffuse reflectance) 2964, 2835, 2483 (w), 2350 (w), 2339 (w), 1956 (w), 1490, 1025, 909, 742 (s), 717, 710 (s), 703 (s), 697 (s), 629, cm⁻¹.

HRMS (EI) calcd for C₂₄H₂₆N₂ 342.2096, found 342.2101.

[α]_D²⁵ = -12° (c 1.00, CH₂Cl₂).

Anal. Calcd for C₂₄H₂₆N₂: C, 84.17; H, 7.65; N, 8.18. Found: C, 84.12; H, 7.64; N, 7.94.

EXAMPLE 7 – Preparation of (2R)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine

The product from Example 6 was dissolved in DMF (150 kg) at 25°C. The solution in the 1200 L reactor was cooled to 20°C to 25°C and transferred into the 1200 L receiver with a DMF rinse (30 kg). Anhydrous K₂CO₃ (103 kg) was added to the 1200 L reactor. The solution in the 1200 L receiver was transferred to the 1200 L reactor with a DMF rinse (30 kg). The 2,3-dichloropyrazine (48.5 kg) was added with a DMF rinse (4 L). The 1200 L reactor was heated to reflux at 127°C to 133°C. Samples were taken every 12-18 h and monitored by GC. The reaction was complete in 41.5 h. The contents of the 1200 L reactor were cooled to 35°C to 45°C and transferred onto a 48" Nutsche filter sending the filtrate sent to the 1200 L receiver. MtBE rinses (2x200 kg at 35°C to 45°C) of the 1200 L reactor were transferred onto the 48" Nutsche filter sending the rinses to the 1200 L receiver. The filtrate in the 1200 L receiver was transferred to the 1200 L reactor with a MtBE rinse (50 kg). The solution in the 1200 L reactor was concentrated under vacuum to remove MtBE and DMF. MgSO₄ was added to a 48" Nutsche (181 kg) and the 1200 L receiver (45 kg). MtBE (625 kg) was added to the 1200 L reactor, heated to 40°C to 45°C, and stirred to dissolve the title compound. The solution was cooled to 15°C to 30°C and transferred into the 1200 L receiver with a MtBE rinse (100 kg). The slurry in the 1200 L receiver was stirred for 3.5 h and filtered onto the 48" Nutsche filter sending the filtrate into the 1200 L reactor. The 1200 L receiver was rinsed with MtBE (2x250 kg at 15°C to 30°C) and transferred to the 48 " Nutsche filter sending the rinses into the 1200 L reactor. The filtrate in the 1200 L reactor was distilled under vacuum to afford the title compound of 87.83% GC purity. The yield was quantitative.

¹H NMR (400 MHz, d₆-DMSO at 87°C): δ 8.18 (1H, d, J = 2.5 Hz), 7.89 (1H, d, J = 2.1 Hz), 7.47 (6H, d, J = 7.6 Hz), 7.31 (6H, s, J = 7.6 Hz), 7.18 (3H, t, J = 7.4 Hz), 4.31-4.26 (1H, m), 3.66 (1H, ddd, J = 12.4, 9.9, 2.5 Hz), 3.48 (1H, br d, J = 12.7 Hz), 2.72 (1H, br d, J = 11.2 Hz), 2.59 (1H, br d, J = 10.7 Hz), 2.49 (d₆-DMSO, reference), 2.14 (1H, br dd, J = 11.2, 2.0 Hz), 1.89 (1H, br t, J = 9.9 Hz), 1.37 (3H, d, J = 6.6 Hz).

¹³C NMR (100 MHz, d₆-DMSO at 87°C): δ 153.85 (s), 141.51 (s), 139.79 (d), 134.86 (d), 128.57 (d), 127.02 (d), 135.60 (d), 76.07 (s), 52.09 (t), 50.84 (d), 47.68 (t), 44.40 (t), 39.52 (d₆-DMSO, reference), 15.39 (q).

IR (diffuse reflectance) 2963 (s), 2350 (w), 2317 (w), 1959 (w), 1921 (w), 1906 (w), 1501, 1488, 1465, 1443 (s), 1411 (s), 1143 (s), 1022, 744, 708 (s), cm⁻¹.

HRMS (FAB) calcd for C₂₈H₂₇ClN₄ + H₁ 455.2002, found 455.2004.

[α]_D²⁵ = -36° (c 0.98, CH₂Cl₂).

Anal. Calcd for C₂₈H₂₇ClN₄: C, 73.91; H, 5.98; N, 12.31; Cl, 7.79. Found: C, 74.26; H, 6.84; N, 10.74.

EXAMPLE 8 – Preparation of (2R)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine

To the 1200 L reactor containing the product from Example 7 was added IPA (707 kg). Stirring at reflux (77°C to 83°C) was continued until dissolution was complete. The solution in the 1200 L reactor was transferred to the 4000 L reactor with a IPA rinse (100 kg) and heated to reflux (77°C to 83°C). A 10% sulfuric acid solution (giving a clean reaction) was prepared in the 4000 L receiver by adding water (245 L) and 98% H₂SO₄ (27 kg). The H₂SO₄ solution was stirred for 10 minutes at 20°C to 60°C and transferred to the 4000 L reactor while maintaining the 4000 L reactor temperature between 60°C to 85°C. The 4000 L receiver was rinsed with water (25 L) sending the rinse to the 4000 L reactor. The 4000 L reactor was heated to reflux (77°C to 83°C). The reaction was monitored by GC and was complete in 30 min. The contents in the 4000 L reactor were cooled to 35°C to 40°C. The IPA was removed under vacuum distillation. Water (1265 L) and MtBE (511 kg) was added to the 4000 L reactor. After stirring and settling, the aqueous layer was transferred into the 4000 L receiver. The MtBE layer in the 4000 L reactor was disposed. MtBE (511 kg) was added to the 4000 L receiver, the solution was stirred and settled for at least 15 min. The aqueous layer was transferred to the 4000 L reactor. The MtBE layer in the 4000 L receiver was disposed. The pH was adjusted to 9.5 to 10.5 using 47% K₂CO₃ (206 kg) with a water rinse (50 L) of the add line. CH₂Cl₂ (2 x 765 L, 1 x 255 L) was used to extract the title compound from the 4000 L reactor sending the CH₂Cl₂ extractions into the 4000 L receiver. The organic solution

in the 4000 L receiver was transferred to the 4000 L reactor with a CH₂Cl₂ rinse (100 L). The CH₂Cl₂ was removed by distillation. Toluene (1000 L) was added to the 4000 L reactor and subsequently removed by vacuum distillation to afford a toluene solution of the title compound of 95.77% GC purity. The yield was quantitative.

Summarizing the steps of Examples 5-8, the yield of the product from Example 8 was 67% starting from (*R*)-2-methylpiperazine, L-tartrate. According to D1, the yield of the same product was 60% starting from (*R*)-2-methylpiperazine, L-tartrate. The step described in D1 corresponding to the process in Example 8 was performed in a mixture of hot ethanol and 10% aqueous hydrochloric acid.

¹H NMR (400 MHz, CDCl₃): δ 8.12 (1H, d, J = 2.6 Hz), 7.87 (1H, d, J = 2.6 Hz), 4.18-4.12 (1H, m), 3.39 (2H, t, J = 3.6 Hz), 3.12 (1H, dd, J = 12.2, 3.6 Hz), 3.06 (1H, dt, J = 12.2, 3.6 Hz), 3.02-2.95 (1H, m), 2.81 (1H, dd, J = 12.2, 4.0 Hz), 1.90 (1H, br s), 1.21 (3H, d, J = 6.6 Hz), 0.00 (TMS, reference).

¹³C NMR (100 MHz, CDCl₃): δ 155.15 (s), 140.85 (s), 139.93 (d), 135.15 (d), 51.06 (d), 50.98 (t), 45.94 (t), 45.10 (t), 14.72 (q), 0.00 (TMS, reference).

IR (liq.) 2940, 2389 (w), 2149 (w), 1996 (w), 1556 (s), 1504 (s), 1459 (s), 1440 (s), 1415 (s), 1367, 1330 (s), 1132 (s), 1107 (s), 1100 (s), 1049 (s), cm⁻¹.

[α]_D²⁵ = -43° (c 0.59, CH₂Cl₂).

HRMS (FAB) calcd for C₉H₁₃ClN₄ + H₁ 213.0907, found 213.0909.

Anal. Calcd for C₉H₁₃ClN₄: C, 50.83; H, 6.16; N, 26.34. Found: C, 50.48; H, 6.19; N, 26.47.

EXAMPLE 9 – Preparation of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine, hydrochloride

To a 200 L reactor was added the product from Example 8 in toluene solution (ca 25 kg thereof in 93 kg solution) with an EtOAc rinse (10 kg). Solvent was removed using vacuum until a volume of 35 L was reached. To remove most of the toluene, EtOAc (2 x 90 kg) was added to the 200 L reactor and distilled until a volume of 35 L was reached. EtOAc (90 kg) was added to the 200 L reactor and heated to reflux (78°C). The slurry in the 200 L reactor was cooled to 55°C to 60°C, filtered through a bag filter to remove solid impurities sending the filtrate to the 200 L receiver. The 200 L reactor was rinsed with EtOAc (5 kg) sending the rinse into the 200 L receiver. The solution in the 200 L receiver was transferred to the 400 L reactor with an EtOAc rinse (5 kg). 37% HCl (9.0 kg) was added to the 200 L receiver and rinsed with EtOAc (9 kg). The 37% HCl solution was transferred to the 200 L

reactor while maintaining a temperature of 73°C to 80°C in the 200 L reactor. The solution in the 200 L reactor was maintained at 72°C for 10 min. The solution was cooled to -25°C to -30°C for 2h 45 min. The resulting slurry in the 200 L reactor was filtered onto an 18"

Nutsche filter sending the filtrate into the 200 L receiver. The reactor and nutsche were rinsed with -25°C EtOAc (3 x 45 kg) sending the filtrate into drums. The filter cake was dried to afford 24.5 kg (79%) of the title compound of 98.4% GC purity. A 2nd similar run gave 23.1 kg for a total of 47.6 kg (77%).

M.Pt. 209.0-210.5°C.

¹H NMR (400 MHz, d₄-MeOH): δ 8.29 (1H, d, J = 2.6 Hz), 8.07 (1H, d, J = 2.6 Hz), 4.83 (2H, br s), 4.30 (1H, sextet, J = 5.8 Hz), 3.68-3.56 (2H, m), 3.40 (1H, dd, J = 14.2, 4.7 Hz), 3.37 (1H, m), 3.31 (1H, br s), 3.23 (1H, dd, J = 12.7, 5.1 Hz), 1.30 (3H, d, J = 6.6 Hz), 0.00 (TMS, reference).

¹³C NMR (100 MHz, d₄-MeOH): δ 155.45 (s), 143.28 (s), 141.90 (d), 138.91 (d), 50.32 (d), 48.60 (t), 44.26 (t), 43.41 (t), 15.41 (q), 0.00 (TMS, reference).

IR (diffuse reflectance) 2934, 2802, 2781 (s), 2744 (s), 2717, 2684, 2470, 2425, 2351 (w), 2335 (w), 2269 (w), 1453 (s), 1412 (s), 1148 (s), 881 cm⁻¹.

HRMS (FAB) calcd for C₉H₁₃ClN₄ + H₁ 213.0907, found 213.0912.

[α]_D²⁵ = -24° (c 0.92, water).

Anal. Calcd for C₉H₁₃ClN₄.HCl: C, 43.39; H, 5.66; N, 22.49. Found: C, 43.48; H, 5.75; N, 22.37.

EXAMPLE 10 – Preparation of (2R)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, fumarate

To a 400 L reactor was added a solution of the product from Example 2A in MeOH (ca. 12.4 kg in 90 L solution) followed by a MeOH rinse (5 kg). The solution was distilled to a volume of 50 L. The fumaric acid (4.6 kg) was added to the 400 L reactor using an addition funnel. The slurry was stirred for 45 min at 25°C to 30°C until all the solids dissolved. MtBE (133 kg) was slowly added, the resulting slurry was cooled to 30°C, and the solids were filtered onto an 18" Nutsche sending the filtrate into the 400 L receiver. The 400 L reactor and 18" Nutsche were rinsed with MtBE (16 kg) sending the rinse into the 400 L receiver. The title compound was dried and packaged to yield 12.3 kg (58%) of 97.4% GC purity. The product purity can be slightly improved by repeating the crystallization.

M.Pt. 125.5-126.5°C.

¹H NMR (400 MHz, d₄-MeOH): δ 8.27 (1H, d, J = 3.1 Hz), 8.15 (1H, d, J = 4.5 Hz), 7.78 (1H, d, J = 2.6 Hz), 7.65 (1H, d, J = 23.1 Hz), 7.48 (1H, dd, J = 8.1, 2.5 Hz), 7.38 (1H, dd, J = 8.5, 4.9 Hz), 6.69 (2H, s), 4.36 (4H, br s), 4.74 (2H, q, J = 4.6 Hz), 4.69-4.63 (1H, m), 4.48 (1H, t, J = 4.3 Hz), 3.99 (1H, dt, J = 11.7, 3.1 Hz), 3.51 (1H, ddd, J = 14.7, 11.2, 3.0 Hz), 3.30-3.26 (2H, m), 3.22-3.15 (2H, m), 1.28 (3H, d, J = 6.7 Hz), 0.00 (TMS, reference).

¹³C NMR (100 MHz, d₄-MeOH): δ 171.20 (s), 156.98 (s), 152.27 (s), 147.16 (s), 142.67 (d), 138.51 (d), 136.14 (d), 135.11 (d), 133.33 (d), 125.94 (d), 123.39 (d), 67.97 (t), 65.75 (t), 48.16 (t), 48.54 (d), 44.17 (t), 40.07 (t), 14.79 (q), 0.00 (TMS, reference).

IR (diffuse reflectance) 3047 (s), 3028 (s,b), 2985 (s,b), 2976 (s), 2962 (s), 2350 (w), 2339 (w), 2318 (w), 2063 (w), 1990 (w), 1710 (s), 1442 (s), 1261 (s), 1206 (s), 1190 (s), cm⁻¹.
[α]_D²⁵ = -20° (c 0.98, water).

Anal. Calcd for C₁₆H₂₁N₅O₂·C₄H₄O₄: C, 55.68; H, 5.84; N, 16.23. Found: C, 55.01; H, 5.88; N, 15.65.

Example 11 - Comparison of the properties of different salts of Example 2A

The following salts were prepared by Chemical development, Stockholm and Kalamazoo and evaluated by Pharmaceutical Development, Nerviano:

- (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine (free base; see Example 2A)
- (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, L-malate (see Example 3A)
- (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, fumarate (see Example 10)
- (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, hydrochloride (see D1, Example 173)
- (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, acetate
- (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, benzoate
- (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, cinnamate.

Identification (NMR and elementary analysis)

Positive data for most salts, except for the free-base and the hydrochloride. NMR analysis was positive for the free-base. The hydrochloride showed problems related to the stoichiometric ratio.

Crystallinity (x-rays)

Positive results for free base, acetate, benzoate, L-malate, and cinnamate were obtained. Most samples were substantially crystalline. Hydrochloride and especially fumarate came out to be of lower crystallinity.

Hygroscopicity (DVS, storage at RT/different RHs and at 40°C/75%RH)

Positive results were obtained on benzoate, L-malate, and cinnamate. They showed lower hygroscopicity ($\leq 5\%$ equilibrium moisture contents at RT/90%RH). Moisture-related effects on the bulk solid state properties of the benzoate salt were observed (cake formation and darkening) but without changes on the DSC pattern. Acetate and hydrochloride were deliquescent (the former at RT/75%RH while the latter at higher values). Fumarate was subjected to hydrate conversion upon exposition at RT/90% RH. Exposition to 40°C/75% RH showed good behavior of cinnamate and L-malate, recovered as powders showing substantially unchanged properties.

Solubility (Water, pH 1.2 and pH 7.4 buffers)

Positive results were obtained for all salts ($\cong 300$ mg/ml FBE) with the exception of cinnamate whose solubility in water is about 8-10 mg/ml FBE.

Intrinsic Dissolution Rate (pH 1.2)

All results confirmed the solubility data. IDRs were about 10^{-1} mgsec⁻¹cm⁻² while cinnamate had much lower dissolution rate (10^{-3} that means -2 order of magnitude lower than the others). Acetate salt showed problem during tablet preparation (sticking to the punch).

Chemical stability (2 weeks at 65°C)

Positive data for benzoate, L-malate, and cinnamate were obtained. Fumarate and acetate seemed to be the less stable salts either at RT/90% RH and at high temperature. Formation of the reaction product with fumaric acid was also observed. These samples showed unchanged DSC patterns except for the acetate (peak broadening).

Polymorphism

The tested conditions (RT evaporation, cooling crystallization at -20°C in different solvent or mixtures) did not show other polymorphs of benzoate and L-malate salts.

Manufacturing method

Benzoate, L-malate, and cinnamate seemed to be easier to crystallize than fumarate, hydrochloride and free base.

5

Toxicological acceptability

There are few marketed drugs that are cinnamate salts and no much information in literature. Oral LD₅₀s in rats are about 2500 mg/kg and 5 g/kg in mice.

Malic acid is an intermediate in the citric acid cycle and occurs naturally in apple and many fruits. It is FDA approved as food additive. A lot of drugs are approved in Europe and US as malate salts.

In summary, of the salts examined above, (2*R*)-methyl-1-{3-[2-(3-pyridinyloxy)ethoxy]-2-pyrazinyl}piperazine, L-malate seems to be the best choice of the salts examined due to its high crystallinity, low hygroscopicity, high chemical stability and low toxicity.

Example 12 - Comparison of the properties of different salts of Example 2B

The following salts have been prepared:

- (2*R*)-1-(3-{2-[(2-ethoxy-3-pyridinyl)oxy]ethoxy}-2-pyrazinyl)-2-methylpiperazine (free base, see Example 2B)
- (2*R*)-1-(3-{2-[(2-ethoxy-3-pyridinyl)oxy]ethoxy}-2-pyrazinyl)-2-methylpiperazine, acetate
- (2*R*)-1-(3-{2-[(2-ethoxy-3-pyridinyl)oxy]ethoxy}-2-pyrazinyl)-2-methylpiperazine, fumarate (see D1, Example 200)
- (2*R*)-1-(3-{2-[(2-ethoxy-3-pyridinyl)oxy]ethoxy}-2-pyrazinyl)-2-methylpiperazine, succinate (see Example 3B).

Crystallinity (powder x-ray diffraction)

For the succinate salt, the degree of crystallinity was very high. The degree of crystallinity decreased in the order succinate>acetate>fumarate.

Thermal properties

For the free base and the acetate salt the melting temperature onset was low (81°C and 96°C, respectively). These low melting points can be drawbacks in future development work.

For the succinate and fumarate salts the thermal properties were fair since the extrapolated melting temperature onset was 123°C and 149°C, respectively.

Hygroscopicity

- 5 All but the acetate salt had acceptable properties with respect to hygroscopicity. The acetate salt was very hygroscopic and showed a pronounced hysteresis at high values of relative humidity.

Solubility and intrinsic dissolution rate

- 10 The solubility of the succinate salt, in all investigate media, was at least 250 mg/ml. For the fumarate salt the solubility was >93 mg/ml in 0.9 % w/w NaCl and >68 mg/ml in phosphate buffer at pH 7.4. Hence, the solubility of succinate was in all instances higher than the fumarate salt.

- The intrinsic dissolution rate studies indicated the dissolution rate to increase
15 according to the order base<fumarate<succinate.

In conclusion, with the respect to the solubility and the intrinsic dissolution rate, the succinate salt proved to have the most favourable qualities of the salts and the free base studied.

Impurities

20 Analysis (LC-UV-MS) of the fumarate salt revealed the presence of a fumarate adduct (~1%) and the formation of dimers of the free base (0.2%). The fumarate adduct may be a Michael adduct between the piperazine ring and fumaric acid.

Conclusion

- 25 The succinate salt hade the overall best state properties, good crystallinity, a relatively high melting point, a high solubility and a high intrinsic dissolution rate. The acetate salt proved to be both hygroscopic and to have a low melting range. The solubility, intrinsic dissolution rate and the degree of crystallinity of the fumarate salt was inferior to that of the
30 succinate salt. In view of the properties of the different salts investigated the best candidate for further development is the succinate salt.

Example 13 - Comparison of the properties of different salts of Example 2C

The following salts have been prepared:

- (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine (free base; see Example 2C)
- (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, phosphoric acid salt (see Example 3C)
- 5 - (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, acetate
- (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, citrate
- (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, edetate
- 10 - (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, oxalate
- (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, succinate
- 15 - (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, D-tartrate
- (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, 1,3,5-benzenetricarboxylate
- (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, galactareate (mucic acid salt)
- 20 - (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, 1,5-naphthalenedisulfonate
- (2*R*)-1-[3-(2-{{2-(2-dimethylaminoethoxy)-3-pyridinyl}oxy}ethoxy)-2-pyrazinyl]-2-methylpiperazine, terephthalate.

25

Crystallinity (powder x-ray diffraction)

The acetate- and the oxalate salt had a high degree of crystallinity whilst the crystallinity of the succinate salt and the phosphoric acid salt ranged from predominantly crystalline to moderately crystallinity. The D-tartrate salt had a low degree of crystallinity.

- 30 Two salts, edetate and citrate proved to be practically amorphous. The free base is a viscous oil at ambient conditions. The benzenetricarboxylate, naphthalenedisulfonate and terephthalate were predominantly crystalline, whilst the galactareate was partially amorphous.

Hygroscopicity

The acetate, citrate, edetate, succinate and D-tartrate salts proved to be very hygroscopic and deliquescent. Also the free base was hygroscopic. The hygroscopicity of the the oxalate and phosphoric acid salt was low. The hygroscopicity of the
5 benzenetricarboxylate, naphthalenedisulfonate and terephthalate was low, moderate and high (deliquescent), respectively.

Solubility


The solubility of the phosphoric acid salt was, in all investigated media, not less than
10 590 mg/ml. Preliminary studies indicated the benzenetricarboxylic acid to have a solubility of >19 mg/ml in SGF (simulated gastric fluid without enzymes; pH of filtrate 2.9), 11 mg/ml in SIF (simulated intestinal fluid without enzymes; pH of filtrate 4.2) and in purified water \approx 4 mg/ml (pH of filtrate 3.6). The solubility of the naphthalenedisulfonate was high >540 mg/ml. Preliminary studies of the solubility of the terephthalate salt in purified water indicated the
15 solubility to be low (<1 mg/ml).

Toxicological acceptability

Regarding the oxalate salt although its solid state properties are good it was deemed inappropriate due to toxicological reasons. The benzenetricarboxylate salt had the overall
20 most promising solid state properties but the solubility was lower than for the phosphoric acid salt and requires a toxicological evaluation. The phosphoric acid salt was deemed to be the best choice of the salts regarding high crystallinity, low hygroscopicity, high solubility, and low toxicity.

Claims


1. A process for the preparation of compounds of the general formula (I):



5


comprising the steps of:

- (i) O-alkylating a 3-pyridinol derivative of the general formula (II) or the corresponding hydrochloride:




10


to give another 3-pyridinol derivative of the general formula (III):



- (ii) reacting the 3-pyridinol derivative of the general formula (III) with (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpyrazine of the formula (IV) in the presence of an alkali metal tert-butoxide or an alkali earth metal tert-butoxide,



to give a compound of the general formula (V):




- (iii) converting the compound of the general formula (V) to the compound of the general formula (I) by treatment with a pharmaceutically acceptable acid of the formula HA,

wherein each of R₁, R₂, R₃, and R₄, independently, is hydrogen, halogen, C₁-C₆-alkyl, C₁-C₆-alkoxy, and di-C₁-C₆-alkylamino-C₂-C₆-alkoxy.


2. The process according to claim 1, wherein

(2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpyrazine of the formula (IV) is prepared by:


(iv) acidification of racemic 2-methylpiperazine of the formula (VI):



5 with L-tartaric acid and fractional crystallization to give (*R*)-2-methylpiperazine, L-tartrate of the formula (VII):




10 (v) basification of (*2R*)-2-methylpiperazine, L-tartrate of the formula (VII) to give (*R*)-2-methylpiperazine of the formula (VIII):




(vi) tritylation of (*R*)-2-methylpiperazine of the formula (VIII) to give (*R*)-3-methyl-1-tritylpiperazine of the formula (IX):

15



- (vii) condensation of (*R*)-3-methyl-1-tritylpiperazine of the formula (IX) with 2,3-dichloropyrazine to give (*2R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X):

5



- (viii) detritylation of (*2R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X) to give (*2R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV),
- (ix) and conversion of (*2R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) to a suitable acid addition salt (e.g, hydrochloride salt).

10

3. The process according to claim 1, wherein the alkali metal tert-butoxide is potassium tert-butoxide.


15

4. The process according to any one of claims 1 to 3, wherein step (ii) is performed in a solvent system including tetrahydrofuran.

5. The process according to any one of claims 1 to 4, wherein step (iv) is performed by using water and ethanol as solvents.


20

6. The process according to any one of claims 1 to 4, wherein step (v) is performed by using a hydroxide as a base.
7. The process according to any one of claims 1 to 4, wherein step (vi) is performed with trityl chloride in the presence of triethylamine.
8. The process according to any one of claims 1 to 4, wherein step (vii) is performed in the presence of an alkali metal carbonate or an alkali earth metal carbonate, with dimethyl formamide as a solvent.
9. The process according to any one of claims 1 to 4, wherein step (viii) is performed in 10% sulfuric acid in isopropanol.
10. A process for the preparation of compounds of the general formula (I):




comprising the steps of:


- (i) reacting a 3-pyridinol derivative of the general formula (II) or the corresponding hydrochloride with ethylene carbonate,




to give another 3-pyridinol derivative of the general formula (III):



- 5 (ii) reacting the 3-pyridinol derivative of the general formula (III) with (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpyrazine of the formula (IV) in the presence of an alkali metal tert-butoxide or an alkali earth metal tert-butoxide,



- 10 to give a compound of the general formula (V):



- (iii) converting the compound of the general formula (V) to the compound of the general formula (I) by treatment with a pharmaceutically acceptable acid of the formula HA,

5


wherein each of R₁, R₂, R₃, and R₄ is hydrogen.

11. The process according to claim 10, wherein

(2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpyrazine of the formula (IV) is prepared by:


10

- (iv) acidification of racemic 2-methylpiperazine of the formula (VI):




with L-tartaric acid and fractional crystallization to give (*R*)-2-methylpiperazine, L-tartrate of the formula (VII):

15




- (v) basification of (2*R*)-2-methylpiperazine, L-tartrate of the formula (VII) to give (*R*)-2-methylpiperazine of the formula (VIII):

5




- (vi) tritylation of (*R*)-2-methylpiperazine of the formula (VIII) to give (*R*)-3-methyl-1-tritylpiperazine of the formula (IX):



10

- (vii) condensation of (*R*)-3-methyl-1-tritylpiperazine of the formula (IX) with 2,3-dichloropyrazine to give (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X):



15

- (viii) detritylation of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X) to give (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV),
- (ix) and conversion of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) to a suitable acid addition salt (e.g. hydrochloride salt).


12. The process according to any one of claims 10 or 11, wherein the alkali metal tert-butoxide is potassium tert-butoxide.

13. The process according to any one of claims 10 to 12, wherein step (ii) is performed in a solvent system including tetrahydrofuran.

14. The process according to any one of claims 10 to 13, wherein


- step (i) is performed in the presence of an alkali metal carbonate or an alkali earth metal carbonate, as a base and dimethylformamide as a solvent,
- step (ii) is performed by reacting the compound of the general formula (III) with the compound of the formula (IV) in the presence of an alkali metal tert-butoxide or an alkali earth metal tert-butoxide, in tetrahydrofuran, or
- the pharmaceutically acceptable acid of the formula HA in step (iii) is L-malic acid.

15. A process for the preparation of compounds of the general formula (I):




comprising the steps of:


- (i) O-alkylating a 3-pyridinol derivative of the general formula (II) or the corresponding hydrochloride,




- 5 to give another 3-pyridinol derivative of the general formula (III):



- (ii) reacting the 3-pyridinol derivative of the general formula (III) with (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpyrazine of the formula (IV) in the presence of potassium tert-butoxide,
- 10



to give a compound of the general formula (V):



- (iii) converting the compound of the general formula (V) to the compound of the general formula (I) by treatment with a pharmaceutically acceptable acid of the formula HA,

5


wherein each of R_1 , R_2 , and R_3 is hydrogen, R_4 is C_1 - C_6 -alkoxy.

16. The process according to claim 15, wherein

(2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpyrazine of the formula (IV) is prepared by:


10

- (iv) acidification of racemic 2-methylpiperazine of the formula (VI):




with L-tartaric acid and fractional crystallization to give (*R*)-2-methylpiperazine, L-tartrate of the formula (VII):

15




- (v) basification of (2*R*)-2-methylpiperazine, L-tartrate of the formula (VII) to give (*R*)-2-methylpiperazine of the formula (VIII):

5




- (vi) tritylation of (*R*)-2-methylpiperazine of the formula (VIII) to give (*R*)-3-methyl-1-tritylpiperazine of the formula (IX):



10

- (vii) condensation of (*R*)-3-methyl-1-tritylpiperazine of the formula (IX) with 2,3-dichloropyrazine to give (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X):



15

- (viii) detritylation of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X) to give (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV),
- (ix) and conversion of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) to a suitable acid addition salt (e.g, hydrochloride salt).


17. The process according to any one of claims 15 or 16, wherein step (ii) is performed in a solvent system including tetrahydrofuran.

18. The process according to any one of claims 15 to 17, wherein R₄ is ethoxy.

19. The process according to any one of claims 15 to 18, wherein


- step (i) is performed by reacting the hydrochloride of the compound of the general formula (II) with 2-chloroethanol with an aqueous solution of a hydroxide,
- step (ii) is performed by reacting the compound of the general formula (III) with the compound of the formula (IV) in the presence of an alkali metal tert-butoxide or an alkali earth metal tert-butoxide, in a solvent system consisting of methyl tert-butylether and tetrahydrofuran, or
- the pharmaceutically acceptable acid of the formula HA in step (iii) is succinic acid.

20. A process for the preparation of compounds of the general formula (I):




comprising the steps of:


- (i) reacting a 3-pyridinol derivative of the general formula (II) or the corresponding hydrochloride with ethylene carbonate,




- 5 to give another 3-pyridinol derivative of the general formula (III):



- (ii) reacting the 3-pyridinol derivative of the general formula (III) with (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpyrazine of the formula (IV) in the presence of an alkali metal tert-butoxide or an alkali earth metal tert-butoxide,
- 10



to give a compound of the general formula (V):



- (iii) converting the compound of the general formula (V) to the compound of the general formula (I) by treatment with a pharmaceutically acceptable acid of the formula HA,

5


wherein each of R₁-R₃ are hydrogen, and R₄ is 2-dimethylaminoethoxy.

21. The process according to claim 20, wherein

(2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpyrazine of the formula (IV) is prepared by:


10

- (iv) acidification of racemic 2-methylpiperazine of the formula (VI):




with L-tartaric acid and fractional crystallization to give (*R*)-2-methylpiperazine, L-tartrate of the formula (VII):

15




- (v) basification of (2*R*)-2-methylpiperazine, L-tartrate of the formula (VII) to give (*R*)-2-methylpiperazine of the formula (VIII):

5




- (vi) tritylation of (*R*)-2-methylpiperazine of the formula (VIII) to give (*R*)-3-methyl-1-tritylpiperazine of the formula (IX):



10

- (vii) condensation of (*R*)-3-methyl-1-tritylpiperazine of the formula (IX) with 2,3-dichloropyrazine to give (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X):



15

- (viii) detritylation of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methyl-4-tritylpiperazine of the formula (X) to give (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV),
- (ix) and conversion of (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) to a suitable acid addition salt (e.g, hydrochloride salt).

22. The process according to any one of claims 20 or 21, wherein the alkali metal tert-butoxide is potassium tert-butoxide.

23. The process according to any one of claims 20 to 22, wherein step (ii) is performed in a solvent system including tetrahydrofuran.

24. The process according to any one of claims 20 to 23, wherein R₄ is 2-dimethylaminoethoxy.

15

25. A process according to any one of claims 20 to 24, wherein


- step (i) is performed in the presence of an alkali metal carbonate or an alkali earth metal carbonate, as a base and dimethyl sulfoxide as a solvent,
- step (ii) is performed by reacting the compound of the general formula (III) with the compound of the formula (IV) in the presence of an alkali metal tert-butoxide or an alkali earth metal tert-butoxide, in tetrahydrofuran, or
- the pharmaceutically acceptable acid of the formula HA in step (iii) is phosphoric acid.

26. A process for preparing (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine, hydrochloride by reacting (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine of the formula (IV) with hydrochloric acid.

27. (2*R*)-1-(3-chloro-2-pyrazinyl)-2-methylpiperazine, hydrochloride.

28. A compound of the formula (I)


54



wherein R_1 - R_4 are hydrogen and HA is L-malic acid.

29. A compound of the formula (I)


5



wherein R_1 - R_3 are hydrogen, R_4 is ethoxy, and HA is succinic acid.

30. A compound of the formula (I)

10



wherein R₁-R₃ are hydrogen, R₄ is 2-dimethylaminoethoxy, and HA is phosphoric acid.

31. A method for treating a serotonin-related disorder, comprising administering to a subject
5 in need thereof an effective amount of a compound according to any one of claims 28 to 30.

32. A method according to claim 31, wherein the serotonin-related disorder is 5-HT_{2C}-
receptor related.

10 33. A method according to any one of claims 31 or 32, wherein the serotonin-related disorder
is selected from obesity and type II diabetes.

34. Use or a compound according to any one of claims 28 to 30 to manufacture a medicament
for the treatment or prophylaxis of a serotonin-related disorder.

15

35. The use according to claim 34, wherein the serotonin-related disorder is 5-HT_{2C}-receptor
related.

20 36. A method according to any one of claims 34 or 35, wherein the serotonin-related disorder
is selected from obesity and type II diabetes.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 03/01043

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 401/14, C07D 401/12, C07D 403/04, A61K 31/444, A61K 31/497, A61P 25/00
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D, A61K, A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 0076984 A2 (PHARMACIA & UPJOHN AB), 21 December 2000 (21.12.00) ----- --	1-36

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 Sept. 2003

Date of mailing of the international search report

25-09-2003

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Eva Johansson/EÖ

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/SE03/01043

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **31-33**
because they relate to subject matter not required to be searched by this Authority, namely:
see next sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/SE03/01043

Claims 31-33 relate to methods of treatment of the human or animal body by surgery or by therapy or diagnostic methods practised on the human or animal body (PCT Rule 39.1(iv)). Nevertheless, a search has been executed for these (this) claim(s). The search has been based on the alleged effects of the compounds or compositions.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 03/01043

Patent document cited in search report			Publication date	Patent family member(s)	Publication date
WO	0076984	A2	21/12/00	AU 1428000	A 22/05/00
				AU 4969000	A 02/01/01
				BR 0010783	A 09/04/02
				CA 2374898	A 21/12/00
				CN 1360583	T 24/07/02
				EP 1140404	A 10/10/01
				EP 1178973	A 13/02/02
				JP 2002528280	T 03/09/02
				JP 2003502317	T 21/01/03
				NO 20015686	A 15/01/02
				SE 9901884	D 00/00/00
				US 6465467	B 15/10/02
				US 2003092694	A 15/05/03
